



ferti·lome®



Fruit Tree Spray

With Neem Py

- Can be used up to day of harvest
- 1 ounce per gallon

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**
See Back Panel for
Precautionary Statements

ACTIVE INGREDIENTS:

Pyrethrins.....	0.25%
Piperonyl Butoxide, Technical.....	2.50%
Clarified Hydrophobic Extract of Neem Oil.....	70.00%
OTHER INGREDIENTS:	27.25%
TOTAL:	100.00%

Net Contents One Pint (16 FL. OZS.)

ferti-lome® Fruit Tree Spray

20M-7-0112 FK

FIRST AID

If On Skin Or Clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If In Eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none">• Move the person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

In the event of a medical or chemical emergency, contact 
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 70051-75-7401
EPA Est. No. 7401-TX-01
Visit Us At: www.fertilome.com
(RV060410)



PEEL FROM CORNER OF BOOKLET



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not expected for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



GENERAL INFORMATION: SHAKE WELL BEFORE AND DURING USE

Multi-purpose insecticide/fungicide/miticide. Kills eggs, larval and adult stages of insects. Prevents fungal attack of plants' tissues. Not for use in food handling areas. **ferti·lome® FRUIT TREE SPRAY** is most effective when applied in early to mid-morning or late afternoon when adult insect pests, such as whiteflies, are normally sedentary on the undersides of the foliage. Use spray solutions within several hours of preparation for maximum effectiveness. For best results maintain constant agitation in spray equipment. For optimal performance, do not mix with cold water or apply under cold ambient temperatures (less than 45°F). If product thickens (freezes) before using, place in a warm room (greater than 45°F) for a few days or in warm water (120°F) for 2 to 8 hours until completely thawed. Do not use partially thawed material. Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment. When using in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale. As with other products that contain oils, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn. Do not add adjuvants (spreaders, stickers, extenders, etc.) to **ferti·lome® FRUIT TREE SPRAY**. Use with care on plants with tender tissue. Do not apply to known spray sensitive plant species such as: impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, red maples, some carnation varieties or similar plants without prior testing.



FUNGICIDE USE

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS, and SHRUBS.

APPLICATION INSTRUCTIONS: **ferti-lome® FRUIT TREE SPRAY** is an effective fungicide for the prevention and control of various fungal diseases including powdery mildew, black spot, brown spot, dollar spot, snow mold, downy mildew, anthracnose, rust, leaf spot, botrytis, needle rust, scab and flower, twig and tip blight. As a preventative, apply **ferti-lome® FRUIT TREE SPRAY** on a 7 to 14 day schedule until the potential for disease development is no longer present. To control disease already present, apply **ferti-lome® FRUIT TREE SPRAY** on a 7 day schedule until disease pressure is eliminated. Then continue spraying on a 14 day schedule to prevent the disease from reoccurring. To prevent rust leaf spot diseases, anthracnose and scab, begin applications at the first sign of spring budding. To prevent powdery mildew, apply in mid-summer or when disease is first detected. For brown patch and dollar spot, begin spraying at first sign of disease on a 7 to 14 day interval. For snow mold, begin spraying on a 14 day interval in the fall before the first snow. Continue applications until disease pressure no longer exists.

MIXING INSTRUCTIONS: Mix **ferti-lome® FRUIT TREE SPRAY** at the rate of 2 tablespoons (1 fluid ounce) per gallon of water. Thoroughly mix solution and spray all plant leaf surfaces (including undersides of leaves) until completely wet. Excessive application is unnecessary and to be avoided.



INSECTICIDE USE

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS, and SHRUBS.

APPLICATION INSTRUCTIONS: Apply **ferti-lome® FRUIT TREE SPRAY** at first sign of insects/mites. **ferti-lome® FRUIT TREE SPRAY** provides for a quicker and more complete control, and acts as an “exciter” to flush insect pests out of hiding and into direct contact with spray residues. For control of aphids, spider mites, scale, mealybugs, whiteflies, beetles, loopers, leafminers, leafrollers, armyworms, webworms, weevils, tent caterpillars, fungus gnats, leafhoppers, cabbageworms, squash bugs, squash vine borer, stink bugs, thrips, tomato hornworm, lace bugs, boxelder bugs, grasshoppers, crickets, ants, fleas, ticks, earwigs, cockroaches, mosquitos, chinch bugs, chiggers and many other insect pests. **ferti-lome® FRUIT TREE SPRAY** is most effective when applied on a 7 to 14 day schedule. For heavy insect populations, spray on a 7 day schedule.

MIXING INSTRUCTIONS: Mix **ferti-lome® FRUIT TREE SPRAY** at the rate of 2 tablespoons (1 fluid ounce) per gallon of water. Thoroughly mix solution and spray all plant leaf surfaces (including undersides of leaves) until completely wet. Excessive application is unnecessary and to be avoided. Frequently agitate solution as you spray.



STORAGE AND DISPOSAL

Pesticide Storage: Do not store diluted solutions for later use.

Pesticide Disposal and Container Handling: If Empty: Non refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Voluntary Purchasing Group, Inc. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

This page left blank intentionally.

This page left blank intentionally.



ferti·lome®

Fruit Tree Spray

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 70051-75-7401

EPA Est. No. 7401-TX-1

Visit Us At: www.fertilome.com

(RV060410)



Get the free mobile app at
<http://gettag.mobi>

