

**DREXEL DIURON 80 HERBICIDE is a state restricted use pesticide and is to be distributed only by licensed pesticide dealers. Only certified applicators or persons under the direct supervision of a certified applicator may apply DREXEL DIURON 80 HERBICIDE.**

SECTION 24(c) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON



*Diuron 80*

Herbicide

**EPA Reg. No. 19713-274  
EPA SLN No. WA-000033**

**This label for Drexel Diuron 80 Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2016.**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.

**DIRECTIONS FOR USE**

Perennial ryegrass grown for seed (established stands) / Control of weeds and seedling grasses (such as annual bluegrass and volunteer ryegrass):

- Apply this product only to well-established, vigorous stands of Perennial ryegrass in the fall at the rate of 1 to 2 pounds per acre. Apply in a minimum of 25 gallons of water per acre for ground application, and a minimum of 5 gallons of water per acre for aerial application. Use sufficient water for thorough coverage of weed foliage.
- For best results, make application at the onset of fall rains and before weeds become established (typically October 1 through November 15). Weeds beyond the 2 to 4 leaf stage will usually not be controlled. Use higher rates within the range listed when treating larger weeds and heavier weed infestation. Weed control may be reduced where straw residues have accumulated on the soil surface. Lack of moisture to activate the herbicide may reduce weed control.

## RESTRICTIONS / PRECAUTIONS

- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Do not make more than one application per year.
- Follow all applicable requirements for spray drift management on the EPA registered label.
- Tank mixtures or sequential treatment with other herbicides may reduce crop tolerance and increase risk of crop injury. When using this product in a tank mix or sequential treatment with other pesticides, do not use the maximum rates listed below unless compatibility and the potential for phytotoxicity have been evaluated.
- Crop tolerance may be reduced and the likelihood of crop injury may increase when crop is under stress caused by weather, diseases and insects.
- For use under this SLN label, do not apply this product through any type of irrigation system.
- Certain uses of diuron may be restricted by a U.S. District Court final order. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/natresources/Buffers.aspx> for information regarding pesticides that are impacted by the final order.

## WSDA CONTAINER DISPOSAL GUIDANCE

Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.



274WASLN-1211(PRGGFS)

The Drexel logo is a registered trademark of Drexel Chemical Company.