



# E-2

SELECTIVE HERBICIDE

## Take Down Tough Weeds with E-2.

Nothing knocks down unsightly weeds like E-2 herbicide. E-2 combines three powerful active ingredients to control more than 240 broadleaf weeds while providing safety for most desirable grasses and reducing the potential for resistant weed development.

### THE POWER OF THREE IN ONE FORMULATION

- Offers the power of three proven herbicides in one convenient pre-mixed formulation
- Excellent control of more than 240 broadleaf weeds including kochia, horsetail, field bindweed, thistles, plantain and other hard-to-control species
- Can be used for selective weed control in desirable grasses and as a tank-mix partner with products like ProClipse™ for cleaning up emerged weeds on bareground areas such as utility substations, industrial areas, highway medians and guardrails
- High-performing weed control in pasture and rangeland with control of tough species like hemp dogbane and dogfennel

**EPA REG. NO.** 228-442

**ACTIVE INGR.** 2,4-D (39.53%),  
fluroxypyr (5.90%),  
dicamba (4.10%)

**FORMULATION** liquid

**CHEM. FAMILY** phenoxyaliphatic acid,  
pyridines, benzoic acids

**GROUP NO.** 4

**MODE OF ACT.** auxin growth regulator

**SIGNAL WORD** warning

**RESTRICTED** no

**PACKAGE SIZE** 2 x 2.5 gal, 30 gal,  
250 gal tote



### KEY USES

Rights-of-way, railroads and roadsides, industrial sites, fence row and farmstead, non-irrigation ditch banks, recreational areas, pasture and rangeland, Conservation Reserve Programs (CRP) Land

### KEY WEEDS CONTROLLED


(See product label for complete list)

Alder  
Beggartick  
Black-eyed Susan  
Black medic  
Buckhorn  
Carolina geranium  
Carpetweed  
Chicory  
Clover  
Dock  
False sunflower  
Garlic mustard  
Ground ivy  
Hemp  
Henbit  
Honeysuckle  
Jimsonweed  
Kochia  
Knotweed  
Lambsquarters  
Marshelder  
Morningglory  
Mouseear chickweed  
Mustard  
Pigweed  
Purslane  
Ragweed  
Shepherd's-purse  
Spatterdock  
Speedwell  
Spurge  
Stinging nettle  
Tansy ragwort  
Thistle  
Tumbleweed  
Vetch  
Virginia creeper  
Wild four-o'clock  
Willow  
Yarrow  
Yellow rocket

### APPLICATION RECOMMENDATIONS

#### WEED CONTROL PERCENTAGES

Field Bindweed Source: Dr. Phil Westra, Colorado State University, 2009  
Buckhorn Plantain Source: P.L. Hipkins and H.L. Witt, Virginia Tech Noncrop and Aquatic Weed Research, 2009  
Wild Carrot Source: P.L. Hipkins and H.L. Witt, Virginia Tech Noncrop and Aquatic Weed Research, 2009



#### NON-CROP

For control of broadleaf weeds, mix 2 to 5 pints per acre in adequate water to thoroughly saturate all weeds with spray mixture. This may require a spray volume of 50 to 300 gallons of water per acre.

#### PASTURES, RANGELAND AND GRASS (HAY, SILAGE)

Some weed species will require tank mixes for adequate control. For these uses, mix 2 to 5 pints per acre. Retreatment may be made as needed; however the maximum seasonal application rate is 2 pounds 2,4-D acid equivalent per acre per application. Minimum of 21 days between applications, limited to 2 applications per year. The pre-harvest interval (PHI) is 7 days (for cut forage for hay).

**Grazing and feeding non-lactating animals:** There is no waiting period between treatment and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

**Grazing and feeding lactating animals:** Do not graze lactating dairy animals within 7 days of treatment.

**Dry hay and silage:** Treated grasses may be harvested for dry hay or silage but do not harvest within 7 days of treatment.

#### CRP AND GENERAL FARMSTEAD

Some weed species require tank mixes for adequate control. For these uses, mix 2 to 5 pints per acre. Retreatment may be made as needed; however the maximum seasonal application rate is 2 pounds 2,4-D acid equivalent per acre per application. For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

See label for complete application rates and recommendations.



More where it matters most.  
[www.nufarm.com/us](http://www.nufarm.com/us)

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit [www.nufarm.com/us](http://www.nufarm.com/us) to download a full label.  
©2013 Nufarm. Important: Always read and follow label instructions. Clean Slate™ and ProClipse™ are trademarks of Nufarm. Garlon® and Milestone® are trademarks of Dow AgroSciences LLC. 13-VM-0007



Scan this code with your smartphone to visit CDMS and view the complete product label.