Prime Part #/Name:	SourceOPC	Revision *	Art File No. <u>0810-013</u> PLEASE CHECK ONE Please resubmit new proof with	
Customer: H	lappy Jack	Proof 10/21/08	 changes noted Approved / Proceed with order 	
PLEASE	E CHECK ONE: Does this label nee	Date:		
	d by:	-	_	
Approval Date	Comments:			
We want ever *If an error *Proofs retu	PLEASE t this proof carefully, including all the co ything right the first time. Please keep t to the copy is approved and printed, it w urned quickly get the earliest press date is returned within 90 days, we must inve	these facts in mind: vill be necessary to char s.	s proof is exactly what will be printed. rge you for the production costs.	
	8 oz. Mange Medicine	Printing Method		
	4.75" X 6.25" - Rectangle R65263 Core Dia: 3		1: <u>Semi-Gloss/C2500</u> 2: 1 Mil Clear Polypropylene	
Chicale		Fhank you.	eSourceOPC 336-661-3300.	Core Dia. = 3"
		Dieline (Does not print)	Label Carrier (Does not print)	4 X CR
	Α			_ = .125"
				= .125"
4.75"	FIRST AD FON SKIN OR CLOTHING - Take off contaminated of plenty of water for 10 - 15 minutes. Call a poison control IF IN EYES - Hold eve open and rinse slowly and ge Remove contact lenses after five minutes then continu- center or doctor for treatment advice. IF SWALLOWED - Call a poison control center or doct sip a glass of water if able to swallow. Do not induce vomi- control center or doctor. Do not give anything by IF INHALED - Move person to fresh air. If person is not center or doctor for treatment advice. Have the product container or label with you when calling going for treatment. PRECAUTIONARY STA KAZARDS TO HUMANS ADD DO Causes moderate eye damage. Harmful if absorbed by individuals. Wash thoroughly with soap and water affect control cothing. Prolonged or repeated use may individuals. Wash thoroughly with soap and water affect cothing and wash before reuse.	center or doctor for treatment advice infly with water for 15 - 20 minutes ue rinsing eye. Call a poison contro tor for treatment advice. Have perso ting unless told to do so by the poiso mouth to an unconscious person b breathing call 911 or an ambulance uth, if possible. Call a poison contro of the poison control center or doctor of TEMENTS DMESTIC ANIMALS through skin. Do not get in eyes, of cause alleroic reactions in som	Provide a second	= .125"
	IF ON SKIN OR CLOTHING - Take off contaminated c plenty of water for 10 - 15 minutes. Call a poison control IF IN EYES - Hold eye open and rinse slowly and ge Remove contact lenses after five minutes then continu center or doctor for treatment advice. IF SWALLOWED - Call a poison control center or doc is of water if able to swallow. Do not induce vomi control center or doctor. Do not give anything by IF INHALED - Move person to fresh air. If person is not then give artificial respiration, preferably mouth to mou center or doctor for treatment advice. Here the product container or label with you when calling going for treatment.	center or doctor for treatment advice infly with water for 15 - 20 minutes ae rinsing eye. Call a poison contro tor for treatment advice. Have perso ting unless told to do so by the poiso mouth to an unconscious person b breathing call 911 or an ambulance uth, if possible. Call a poison contro to the poison control center or doctor of the poison control center of doctor of the poison center of docto	nn Bad SARCOPTIC MANGE MEDICINE	= .125"
	IF ON SKIN OR CLOTHING - Take off contaminated c plenty of water for 10 - 15 minutes. Call a poison control IF IN EYES - Hold eye open and rinse slowly and ge Remove contact lenses after five minutes then continu center or doctor for treatment advice. IF SWALLOWED - Call a poison control center or doc sip a glass of water if able to swallow. Do not induce vomi control center or doctor. Do not give anything by IF INHALED - Move person to fresh air. If person is not then give artificial respiration, preferably mouth to mou- center or doctor for treatment advice. Have the product container or label with you when calling going for treatment.	center or doctor for treatment advice netwith with water for 15 - 20 minutes ue rinsing eye. Call a poison contro tor for treatment advice. Have perso ting unless told to do so by the poiso mouth to an unconscious person to the treathing call 911 or an ambulance uth, if possible. Call a poison contro the poison control center or doctor of the poison control center of doctor the poison control center of doctor the poison control center or doctor of the poison control center or doctor of the poison control center or doctor of the poison control center of doctor the poison control center of doctor the poison control center of doctor the poison control center of doctor of the poison control center of doctor the poison control center of doctor of solvely, use twice da PORTANT for best results, disinfer foldic check of infection and sour	And	
4.75"	IF ON SKIN OR CLOTHING - Take off contaminated c plenty of water for 10 - 15 minutes. Call a poison control IF IN EYES - Hold eye open and rinse slowly and ge Remove contact lenses after five minutes then continu center or doctor for treatment advice. IF SWALLOWED - Call a poison control center or doc sip a glass of water if able to swallow. Do not induce vomi control center or doctor. Do not give anything by IF INHALED - Move person to fresh air. If person is not then give artificial respiration, preferably mouth to mou- center or doctor for treatment advice. Have the product container or label with you when calling going for treatment.	center or doctor for treatment advice netwith with water for 15 - 20 minutes ar insing eye. Call a poison contro tor for treatment advice. Have perso ting unless told to do so by the poiso mouth to an unconscious person t breathing call 911 or an ambulance the poison control center or doctor of TEMENTS DMESTIC ANIMALS through skin. Do not get in eyes, of cause allergic reactions in som ter handling. Remove contaminate POSAL posal. reuse or refill this container. Place i 0-CLEANUP for disposal instructions an children. So manner inconsistent with its labelin E USING nsult a veterinarian before using th as. Sensitivities may occur after usis cur wash your get with mild scap an consult a veterinarian before using th as. Sensitivities may occur after usis cur wash your goves and work w s that respond slowly, use twice da PORTANT for best results, disinfe riodic check of infection and sour rinarian. Manufactured by: HAPPY JACK, INC. P.O. Box 475/2122 Hwy. 258 1-NC-1 Snow Hill, NC 28580	And	