

# Liqui-Cop® RTS

## COPPER FUNGICIDE GARDEN SPRAY

Controls Many Diseases Attacking  
Citrus and Fruit & Nut Trees

Ready-To-Spray Bottle  
Just Attach to Garden Hose & Spray

Delivers 4 Tsp. of LIQUI-COP RTS per Gallon of Water  
Makes up to 48 Gallons of Spray Solution

### FOR RESIDENTIAL USE ONLY

#### ACTIVE INGREDIENT:

Copper diammonia diacetate complex\*  
[Bis (acetate-O) Diamminocopper] ..... 27.15%  
CAS No. 13822-80-5  
OTHER INGREDIENTS: ..... 72.85%  
TOTAL: ..... 100.00%

\* Metallic Copper Equivalent, 8.0%  
Contains 0.196 lbs. Copper per quart

EPA Reg. No. 54705-7

EPA Est. No. 48498-CA-1

**NET CONTENTS: 1 QUART / 32 FL OZ / 946 mL**



Manufactured for:

**LAWN AND GARDEN PRODUCTS, INC.**

P. O. Box 35000 • Fresno, CA 93745 • (559) 499-2100  
info@montereylawngarden.com

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes eye and skin irritation. Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wear protective eyewear (goggles, face shield or safety glasses), long-sleeved shirt and long pants, shoes and socks and chemical-resistant gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter the treated area until sprays have dried.

The control of fungal disease is based on PREVENTION. Copper pesticides are considered preventative, not curative of plant diseases. Plant surfaces must be completely covered with fungicide to successfully prevent infection.

Seek advice on identifying specific disease problems and solutions from your local Agricultural Extension Agent or other knowledgeable persons.

### INSTRUCTIONS

SHAKE WELL BEFORE USING.

**APPLICATION:** Thoroughly spray tree or bush to the point of runoff, including upper and lower surfaces of any foliage. Do not overspray. Do not apply at less than 7 day intervals.

**NOTE:** This product may be reactive on metal and masonry surfaces such as galvanized roofing. AVOID contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.



**HOW TO USE THIS SPRAYER:** Shake Container Well Before Getting Started.

The sprayer attached to the container is ready-to-use. Simply attach to your garden hose and follow these instructions.

1. Connect sprayer to hose.
2. Turn on water.
3. To begin spraying, point nozzle in the direction you want to spray
4. Bend plastic tab back and turn knob to ON position.
5. Spray evenly to area to be treated.

NOTE: LIQUI-COP RTS is a dark blue liquid. When the spray turns clear it is time to change to a new bottle of product.

6. To stop spraying, turn knob to OFF position.
7. Turn off water.
8. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
9. Disconnect sprayer from hose.



## FRUITS AND NUTS

FRUITS AND NUTS	DISEASE CONTROLLED	INSTRUCTIONS
ALMONDS	Brown Rot	Apply at delayed dormant* bud swell stage.
	Shot Hole	Apply at leaf fall to protect buds and shoots from infection during rainy periods. Reapply every 3 to 4 weeks up until late bud swell (the period in Spring before buds begin to open). Do not apply after full bloom.
APPLES	Apple Scab (Black Spot) Bacterial Canker Blossom and Shoot Blast	Apply post-harvest before fall rains.
APRICOTS	Brown Rot Blossom Blight	Apply when buds show red color until brown bud jackets have fallen.
	Shot Hole	Apply in late fall or early winter before winter rains or snow. Repeat in late winter or early spring before buds break. Do not apply after full bloom.
AVOCADOS	Anthrachnose	Begin application when the flower buds begin to swell and continue at monthly intervals until August.
BLUEBERRIES	Bacterial Canker	Apply before fall rains and again 4 weeks later.
	Cane Canker	Apply before fall rains and again 4 weeks later. In the spring, during wet weather, apply at 10-14 day intervals beginning at leaf emergence.
CANE BERRIES	Anthrachnose Bacterial Blight Leaf and Cane Spot Purple Blotch Yellow Rust	Apply in the fall after harvest.
CHERRIES	Brown Rot Blossom Blight	Apply as a full cover spray at popcorn stage and at full bloom.
CITRUS	Greasy Spot Melanose Pink Pitting Scab	Apply as pre-bloom and post bloom sprays.
CURRANTS, GOOSEBERRIES	Anthrachnose Leaf Spot (Cane Blight)	Make 3 applications starting after harvest, before bloom and after petal fall.
LIMES	Greasy Spot	Apply in June and continue at monthly intervals through August.
MANGOS (Not for use in California)	Anthrachnose	Apply at 30 day intervals from the time the flowers are 2 inches in length until all fruits are set.
OLIVES	Peacock Spot	Make first application before winter rains fall. A second application should be made in early spring if disease is severe.
PEACHES NECTARINES	Bacterial Spot	Apply as a dormant spray.
	Blossom Brown Rot	Apply as a dormant or delayed dormant* spray. Do not apply at or after full bloom.
	Leaf Curl Shot Hole	Apply at leaf fall to protect buds and shoots from infection during rainy periods. Reapply up until late bud swell. Do not apply after full bloom.
PEARS	Blossom Blast	Apply as a dormant spray. Apply only at bud break to control primary infection.
PECANS	Shuck and Kernel Rot Zonate Leaf Spot	For suppression, apply at 2 – 4 week intervals starting at kernel growth and continuing until shucks open. Use shorter interval if frequent rainfall occurs.
PISTACHIOS	Alternaria Late	Apply at 50% and full bloom

	Blight	followed by up to 3 applications at 30-day intervals.
	Botrytis Blight Botryosphaeria Panicle Shoot Blight Septoria Leaf Blight	Make initial application at bud swell and repeat on a 14 – 28 day schedule as dictated by disease conditions. If disease conditions are severe, use the higher rate and shorter spray interval.
PLUMS PRUNES	Bacterial Blast Bacterial Canker Shot Hole	Apply as a dormant spray and at early green bud to full popcorn stage.
WALNUTS	Walnut Blight	Make first application at early pre-bloom. Make second application at late pre-bloom. Make additional applications if disease conditions persist at 7 day intervals.

\* Delayed Dormant – The period just before blossom buds break open.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a secure, locked area away from pets and out of the reach of children. Store only in original container and place in a locked storage area.

**PESTICIDE DISPOSAL/CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The Directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of the product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use of application, all of which are beyond the control of Lawn and Garden Products, Inc. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Lawn and Garden Products, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Lawn and Garden Products, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lawn and Garden Products, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Lawn and Garden Products, Inc.'s election, the replacement of product.

(LG-1112N) 0808/1112(03)