

# Supplemental Label



**For disease control and plant health in the commercial production of greenhouse grown tomatoes and tomato transplants for the home consumer market**

**This label expires on December 31, 2017 and must not be used or distributed after this date.**

**Active Ingredients:**

pyraclostrobin*: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1 <i>H</i> -pyrazol-3-yl]oxy]methyl]phenyl] methoxy-, methyl ester) .....	12.8%
boscalid**: 3-pyridinecarboxamide,2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- .....	25.2%
<b>Other Ingredients:</b> .....	62.0%
<b>Total:</b> .....	100.0%

\* 0.128 oz (0.008 lb) of pyraclostrobin in 1 oz of product  
 \*\* 0.252 oz (0.0158 lb) of boscalid in 1 oz of product

**EPA Reg. No. 7969-251**

---

### Directions For Use

---

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Pageant® Intrinsic™ brand fungicide** container label, EPA Reg. No. 7969-251, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Pageant Intrinsic** before applying.
- Use of **Pageant Intrinsic** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Pageant Intrinsic**.

---

### Application Information

---

Refer to the **Pageant Intrinsic** container label for additional application instructions and restrictions.

## Application Instructions

### Commercial Production of Greenhouse Grown Tomatoes

Apply **Pageant® Intrinsic™ brand fungicide** for control of Botrytis gray mold in the commercial production of greenhouse grown tomatoes.

- Apply **Pageant Intrinsic** as a foliar spray in a minimum of 100 gal/A water volume.
- Allow at least 7 days between applications.
- Applicators and other handlers must wear a NIOSH approved respirator with any N, R, P, or HE filter or NIOSH approval number prefix TC-84A when applying with a mechanically pressurized handgun to greenhouse tomato.
- **DO NOT** use **Pageant Intrinsic** for tomato transplants that are intended for agricultural production fields.
- **DO NOT** tank mix **Pageant Intrinsic** with adjuvants or other agricultural products in commercial production of greenhouse grown tomatoes.
- **DO NOT** apply more than a total of 1.1 lbs ai/A (boscalid + pyraclostrobin) [69 ozs/A of **Pageant Intrinsic**] per year.

**Table 1. Pageant Intrinsic Crop-specific Requirements – Commercial Production of Greenhouse Grown Tomatoes**

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Crop Cycle	Maximum Rate per Crop Cycle (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Commercial production of greenhouse grown tomatoes	Botrytis gray mold <i>Botrytis cinerea</i>	23	3	69	0

**Application Directions.** For control of Botrytis gray mold, apply 23 ozs/A of **Pageant Intrinsic** prior to onset of disease development when conditions favor disease development. **DO NOT** apply more than 69 ozs/A of product per crop cycle.

**Resistance Management.** To limit potential for development of resistance, **DO NOT** make more than three (3) applications of **Pageant Intrinsic** or other **Group 7** or **Group 11** fungicides per crop cycle.

**DO NOT** make more than one (1) application of **Pageant Intrinsic** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

## Tomato Transplants for the Home Consumer Market

Apply **Pageant<sup>®</sup> Intrinsic<sup>™</sup> brand fungicide** for disease control on tomato transplants grown for the home consumer market in greenhouses, lathhouses, or other production structures to control foliar, stem, crown and root diseases listed in **Table 2. Pageant<sup>®</sup> Intrinsic<sup>™</sup> brand fungicide Crop-specific Requirements - Tomato Transplants for the Home Consumer Market.**

**Pageant Intrinsic is not for use on vegetable transplants intended for agricultural production fields.**

To maximize disease control, apply **Pageant Intrinsic** in a regularly scheduled protective spray program and use in a rotation with other fungicides that are not in **Group 7** and **Group 11** fungicides. Because of its high specific activity, **Pageant Intrinsic** has good residual activity against target fungi.

- **DO NOT** exceed 100 gallons per acre spray volume.
- Allow at least 7 days between applications.
- **DO NOT** tank mix adjuvants or other agricultural products with **Pageant Intrinsic** for use on tomatoes.
- **DO NOT** apply more than two (2) consecutive **Pageant Intrinsic** applications in any crop production cycle. Rotate to a fungicide of a different mode of action group (**non-Group 7** or **non-Group 11** fungicides) before reapplying **Pageant Intrinsic**.
- **DO NOT** apply more than three (3) **Pageant Intrinsic** applications to any crop during a growing cycle.
- **DO NOT** apply **Pageant Intrinsic** to consecutive tomato crops within the same production structure. Alternate to other effective fungicides with different modes of action from **Group 7** or **Group 11** fungicides.
- **DO NOT** use **Pageant Intrinsic** for tomato transplants that are intended for agricultural production fields.
- **DO NOT** apply more than a total of 1.9 lbs ai (boscalid + pyraclostrobin) [7.3 lbs (118 ozs) of **Pageant Intrinsic**] in the same production structure per year.

**Table 2. Pageant® Intrinsic™ brand fungicide Crop-specific Requirements – Tomato Transplants for the Home Consumer Market**

<b>Crop</b>	<b>Target Disease</b>	<b>Product Use Rate per Application</b> (ozs/100 gallons)	<b>Application Interval</b> (days)*	<b>Comments</b>
<b>Tomato transplants for the home consumer market</b>	<b>Crown and basal rot</b> <i>Fusarium</i> spp. <i>Rhizoctonia solani</i> <i>Sclerotinia</i> spp.	12 to 18	7 to 14	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. The crown and base of the plant and the soil or potting medium surrounding the crown must be thoroughly covered.
	<b>Damping-off</b> <i>Rhizoctonia</i> spp. <i>Pythium</i> spp.	12 to 18	7 to 14	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. The crown and base of the plant and the soil or potting medium surrounding the crown must be thoroughly covered.
	<b>Downy mildew</b> <i>Peronospora</i> spp. <i>Plasmopara</i> spp.	12 to 18	7 to 10	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
	<b>Spots and blights</b> <i>Alternaria</i> spp. <i>Cercospora</i> spp. <i>Phoma</i> spp. <i>Septoria</i> spp.	8 to 12	7 to 14	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to or at the first disease symptom development.
	<b>Phytophthora blight</b> <i>Phytophthora</i> spp.	18	7 to 10	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to or at the first disease symptom development. Alternate and/or tank mix with chlorothalonil, mancozeb or fixed copper fungicides for late blight protection of tomato.
	<b>Powdery mildew</b> <i>Leveillula</i> spp. <i>Oidiopsis</i> spp.	12	7 to 10	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to or at the first disease symptom development.
	<i>Oidium</i> spp.	6 to 12		
	<b>Rots and blights</b> Botrytis rot <i>Botrytis</i> spp.	12 to 18	7 to 10	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.

**Application Directions.** Apply **Pageant Intrinsic** preventively for production of tomato transplants in greenhouses, lathhouses, or other production structures for home consumer market only.

Begin when conditions are favorable for fungal infection, prior to disease symptom development.

For control of listed diseases, apply **Pageant Intrinsic** as a broadcast or directed spray applied in sufficient water to obtain coverage of the plant crown and plant stem through wetting of the soil surface.

\*The stated interval applies to conditions under which moderate-to-high disease pressure is expected. If conditions are unfavorable for infection, or if disease pressure is absent, the interval can be extended up to 28 days.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Pageant* is a registered trademark of BASF.

*Intrinsic* is a trademark of BASF.

©2014 BASF Corporation  
All rights reserved.

007969-00251.20140717.NVA 2014-04-280-0195

Based on: NVA 2014-04-280-0194

Supersedes: NVA 2014-04-280-0162

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company