

TERRAGUARD® SC

Ornamental Fungicide



AN ORNAMENTAL FUNGICIDE FOR CONTROL OF RHIZOCTONIA ROOT ROT AND OTHER HARD TO CONTROL DISEASES SUCH AS THIELAVIOPSIS, CYLINDROCLADIUM, POWDERY MILDEW, RUST, SCAB AND LEAF SPOTS.

Product Information Bulletin

945

GENERAL INFORMATION

Terraguard® SC is a locally systemic fungicide that provides protection of hard to control diseases. The active ingredient in **Terraguard SC** prevents diseases from attacking leaf surfaces as it penetrates the plant to stop disease development.

Terraguard SC is a 42.14% flowable formulation of the active ingredient Triflumizole

Common Name: Triflumizole

Class: Imidazole

GROUP	3	FUNGICIDE
-------	---	-----------

Packaging: 1 quart bottle, 12 per case

RESTRICTED ENTRY INTERVAL (REI) AND SIGNAL WORD

REI = **12 hours**

Signal Word = **CAUTION**

PERSONAL PROTECTION EQUIPMENT (PPE)

- long sleeved shirt and long pants
- chemical-resistant gloves made of any waterproof material
- shoes plus socks

MODE OF ACTION

Terraguard SC works by inhibiting sterol synthesis. Optimal performance is obtained as a preventative application. **Terraguard SC** can also be applied after disease symptoms appear and provide good eradicant control.

USE SITES

Terraguard SC may be used on ornamental plants grown in commercial greenhouses, shadehouses, interiorscapes, and nurseries, including Christmas tree / conifer plantations.

APPLICATION RATES

Terraguard SC may be applied as a spray, drench, cutting soak or through chemigation equipment.

Apply 2- 8 -16 fluid ounces per 100 gallons depending on disease and application method. See label for specific instructions.

Note:

Cultivars of impatiens have shown sensitivity to applications of **Terraguard SC**. Do not exceed 2 fluid ounces per 100 gallons used on impatiens transplants or plugs.

For optimum results from drench applications, do not irrigate crop with additional water until 24 hours after application.

TANK MIXING

The **Terraguard SC** user is advised to conduct a small scale test for any tank mix compatibility with other products before using on a large scale.

KEY DISEASES CONTROLLED

- Rhizoctonia root rot
- Thielaviopsis
- Cylindrocladium
- Powdery Mildew
- Rusts
- Leaf Spots
- Scab
- Fusarium wilt

RESISTANCE MANAGEMENT: Use the following chart to develop a resistance management strategy. ✓ = labeled

TRADE NAME	Aliette® WDG	FenStop™	Compass® O	OHP Chipco® 26019 N/G OHP 26 GT®-O	OHP 6672™	Strike® 50 WDG	Terraclor® 400 SC	Terraclor® 75% WP	Terraguard® SC	Terrazole® 35% WP	Terrazole® CA	Triact® 70
CLASS	Phosphonate	Imidazolinone	Strobilurin	Dicarbonylimide	Benzimidazoles	Triazole	Aromatic Hydrocarbon	Aromatic Hydrocarbon	Imidazole	Aromatic Hydrocarbons	Aromatic Hydrocarbons	Oil
PATHOGEN												
Alternaria		✓		✓					✓			
Anthracoze		✓			✓							✓
Black Spot		✓			✓							✓
Botrytis		✓		✓	✓				✓			
Botrytis Storage Rot				✓			✓					
Cercospora Leaf Spot					✓	✓						
Cylindrocladium			✓	✓	✓				✓			
Downy Mildew	✓	✓										
Entomosporium Leaf Spot					✓	✓						
Fusarium				✓	✓				✓			
Leaf Spot		✓		✓	✓	✓			✓			
Myrothecium Leaf Spot		✓			✓							
Ovulinia Flower Blight					✓	✓						
Pythium	✓	✓								✓	✓	
Phytophthora Root Rot	✓	✓								✓	✓	
Phytophthora Aerial	✓	✓										
Powdery Mildew		✓			✓	✓			✓			✓
Rhizoctonia		✓		✓	✓		✓		✓			
Rust		✓				✓			✓			✓
Scab		✓			✓				✓			
Sclerotinia					✓	✓						
Septoria Leaf Spot		✓			✓							
Thielaviopsis		✓			✓				✓			
Tip Blight					✓	✓						

Aliette, Compass, 26 GT, and Chipco are registered trademarks of Bayer. FenStop is a trademark of Bayer. Triact is a registered trademark of Ceris USA, LLC. Terraguard and Terraclor are registered trademarks of Chemtura Corporation. Terraclor is a registered trademark of Amvac Chemical Corporation. Strike is a registered trademark of OHP, Inc. OHP 6672 is a trademark of OHP, Inc.

Read label directions and cautions before use.

© OHP, Inc. 06/2008

PO Box 230, Mainland, PA 19451
 Technical Service: 800-356-4647
 www.ohp.com

