

WEED BEATER ULTRA READY TO USE

- **PERFORMS IN BOTH WARM AND COOL WEATHER**
- **WON'T HARM LAWN GRASS WHEN USED AS DIRECTED**
- **FOR NORTHERN AND SOUTHERN GRASSES**
- **SEE RESULTS IN HOURS**
- **RESEED IN 2 WEEKS**

EPA Reg. No. 2217-865-4
EPA Est. No. 4-NY-1

©Bonide Products, Inc. All Rights Reserved

Distributed by
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Precautions and Restrictions:

- Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

PRODUCT FACTS

This product contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

This product offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in turfgrass, e.g. spurge, pennywort (dollarweed), dandelion, and white clover.
- Injury to weeds can be noticed within hours of the application and plant death can occur within 7-14 days.

WHERE TO USE

On Lawns:

Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as: Kentucky bluegrass, Perennial ryegrass, Tall fescue, Red or fine leaf fescues, Mixtures of cool season species, Common bermudagrass, Hybrid bermudagrass, St. Augustinegrass (excluding 'Floratam'), Zoysiagrass.

Prohibitions:

- Do not apply this product to bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not apply to 'Floratam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.

APPLICATION SCHEDULES (WHEN TO APPLY)

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 2 to 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.

APPLICATION SCHEDULES (WHEN TO APPLY) Cont.

- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant bermudagrass, zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.
- Do not apply more than 1.5 lb ai (MCPA)/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

HOW TO USE

For Containers with Trigger Sprayers: This product is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

BROADLEAF WEEDS CONTROLLED

This product will control or suppress the following broadleaf weeds.

BROADLEAF WEEDS

Annual yellow sweetclover, Aster, Austrian fieldcress, Bedstraw, Beggarticks, Betony Florida, Bindweed field, Bird vetch, Bitter wintercress, Bittercress, hairy, Bitterweed, Black-eyed Susan, Black medic, Black Mustard, Blackseed plantain, Blessed thistle, Bloodflower milkweed, Blue lettuce, Blue vervain, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull thistle, Bullnettle, Burclover, Burdock, Burning nettle, Burweed, Buttercup, Buttonweed, Canada thistle, Carolina geranium, Carpetweed, Catchweed Bedstraw, Catnip, Catsear, Chickweed common, Chickweed mouseear, Chicory, Cinquefoil, Clover crimson, Clover hop, Clover red, Clover strawberry, Clover sweet, Clover white, Cockle, Cocklebur, Common mullein, Creeping Jenny (Creeping Charlie), Cudweed, Curly dock, Daisy English, Daisy fleabane, Daisy oxeye, Dandelion, Dichondra, Dogbane, Dogfennel, Dollarweed, Elderberry, False dandelion, Falseflax, False sunflower, Fiddleneck, Florida pusley, Frenchweed, Galinsoga, Goathead, Goldenrod, Ground ivy, Gumweed, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Heathaster, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk vetch, Morningglory, Mouseear hawkweed, Mugwort, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oriental cocklebur, Oxalis, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis (tickseed), Plantain, Poison ivy, Poison oak, Pokeweed,

BROADLEAF WEEDS Cont.

Poorjoe, Prairie sunflower, Prickly lettuce, Prickly sida, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncturevine, Purslane common, Ragweed, Red sorrel, Redroot pigweed, Redstem filaree, Rough cinquefoil, Rough fleabane, Russian pigweed, Russian thistle, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smooth dock, Smooth pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny amaranth, Spiny cocklebur, Spiny sowthistle, Spotted catsear, Spotted spurge, Spurweed, Stinging nettle, Strawberry India mock, Tall nettle, Tall vervain, Tansy ragwort, Tansy mustard, Tanweed, Thistle, Trailing crownvetch, Tumble mustard, Tumble pigweed, Velvetleaf, Venice mallow, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Water Pennywort, Wavyleaf bullthistle, Western clematis, Western salsify, White mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild mustard, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Woodsorrel, Woolly croton, Woolly morningglory, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed and other broadleaf weeds.

STORAGE & DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

CONTAINER DISPOSAL: *If empty:* Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled:* Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment washwater or rinsate.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.