

# CHICKWEED CLOVER & OXALIS KILLER READY-TO-USE

By Isomer Specific Method, Equivalent to:

- \*2-Methyl-4-Chlorophenoxyacetic Acid ..... 0.604%, 0.057 lbs./gal.
- \*\*3,5,6-Trichloro-2-Pyridinyloxyacetic Acid ..0.060%, 0.006 lbs./gal.
- \*\*\*3,6-Dichloro-o-Anisic Acid .....0.060%, 0.006 lbs./gal.

- **KILLS 150+ WEED TYPES, ROOTS AND ALL.**
- **CONTAINS MCPA, TRICLOPYR AND DICAMBA.**
- **BEGINS WORKING OVERNIGHT, RAINPROOF IN HOURS.**
- **CONSISTENTLY FAST CONTROL ON TOUGH WEEDS  
LIKE CLOVER, PLANTAIN, GROUND IVY (CREEPING  
CHARLIE) AND SPURGE.**

Store and transport in an upright position.

EPA Est. No. 4-NY-1

EPA Reg. No. 228-336-4

Buyers Guarantee limited to label claims.

Distributed by  
Bonide Products, Inc.  
6301 Sutliff Road  
Oriskany, NY 13424

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Residential Ornamental Turf Lawns. The maximum number of broadcast applications per treatment site is two per year.

**KILLS:** Poison Ivy, Poison Oak, Alder, Ash, Birch, Beech, Kudzu, Blackberry, Raspberry, Salmonberry, Thimbleberry, Willow, Virginia Creeper, Trumpet Creeper, Vine Maple, Choke Cherry, Canada Thistle, Cocklebur, Ground Ivy, Stinging Nettle, Thistle, Whitebrush, Yellow starthistle and many other broadleaf weeds.

## HOW TO USE

Trigger Spray – Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. One application should be sufficient. Some hard-to-kill broadleaf weeds may require a second treatment in 3 to 4 weeks.

## USE PRECAUTIONS

Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray roots of desirable trees and plants. Maximum kill of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. Do not use on Carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. Do not apply to newly seeded grasses until they have been mowed 3 times.

Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use.

If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

**Entry Restrictions:** Do not allow people or pets to enter the treated area until dusts have settled.

**Application Restrictions:** Avoid contact with eyes, skin, or clothing during and after application. Do not apply this product in a way that will contact persons or pets either directly or through drift. Keep people and pets out of the area during application.

## THIS PRODUCT CONTROLS

Alfalfa, Amaranth, Beadstraw, Bedstraw, Beggartick, Beggarweed, Bindweed, Bitter sneezeweed, Black medic, Bladder campton, Bracken fern, Broomweed, Buckhorn, Buckwheat, Buffalobur, Burclover, Burcucumber, Burdock, Bursage, Buttercup, Canada thistle, Carpetweed, Catnip, Catsear, Chamise, Chamomile, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Coffeeweed, Cornflower, Cornspeedwell, Croton, Daisy, Dandelion, Dock, Dog fennel, Dragonhead mint, Evening primrose, Field pennycress, Fleabane, Florida pusley, Frenchweed, Galinsoga, Geranium, Goatsbeard, Goldenrod, Goosefoot, Gromwell, Ground ivy, Halfshrub sundrop, Hawkweed, Healall, Heartleaf drymary, Hemp sesbania, Hempnettle, Henbit, Horsenettle, Horseweed, Ironweed, Jerusalem artichoke, Jimsonweed, Knapweed, Knapwood, Knawel, Knotweed, Kochia, Ladysthumb, Lambsquarter, Lespedeza, Little starwort, Mallow, Marc's Tail, Marshelder, Matchweed, Mayweed, Milkweed, Morningglory, Mustard, Nightflowering catchfly, Nightshade, Oxalis (Stricta and corniculata), Parsley-piert, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Prickly sida, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red sorrel, Red clover, Redevine, Redstem, Rubberweed, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Snakeweed, Sowthistle, Speedwell, Spiderwort, Spikeweed, Spurge, Spurry, Spurweed, Starbur, Stinging nettle, Stitchwort, Stinkweed, Sumpweed, Sunflower, Tansy ragwort, Teasel, Teaweed, Texas blueweed, Thistle, Toadflax, Trumpet creeper, Velvetleaf, Veronica, Vetch, Waterhemlock, Waterhemp, Whitebrush, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild violet, Wood sorrel, Wormwood, Yankeeweed, Yarrow, Yellow starthistle, Yellow rocket, and many other broadleaf weeds.

## STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep from freezing. To be stored in original container and placed in an area inaccessible to children.

**PESTICIDE DISPOSAL:** *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – If product cannot be used as directed, call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

## FIRST AID

<b>IF: SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Remove and wash contaminated clothing before reuse.

**USER SAFETY RECOMMENDATIONS**

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour washwater on the ground; spray or drain over a large area away from wells or other water sources. Do not contaminate water used for irrigation or domestic purposes.

Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Desirable plant injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Bonide or Seller. All such Risks shall be assumed by Buyer and User, and Buyer and User agree to hold Bonide and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. BONIDE MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT.** Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Bonide, and Buyer assumes the risk of any such use.

To the extent allowed by law, Bonide or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BONIDE AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF BONIDE OR SELLER, THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.