

PRODUCT FACTS

What it Does:

Helps protect your home from invading Ants, except Carpenter and Pharaoh Ants. Also Kills Crickets, Fire Ants, Chinch Bugs, Armyworms (Fall and Yellow-striped), Billbugs, Cutworms, Earwigs, Fleas, Roaches (German & American), Grubs, Sod webworms, European Crane Flies, Ticks and other listed Insects

Where to Use:

Outdoors, on Lawns and as a Band Treatment around House Foundation
For outdoor use around the home only.


Questions & Comments:

Call 1-800-536-8231 or visit our website at www.bonide.com

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Keep children and pets off treated area until product has been watered in and lawn is dry.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

Three Easy Steps to Use this Product

- Step #1: Determine size of the job, measure the area to be treated.
- Step #2: Use spreader setting chart to determine appropriate setting for your spreader.
- Step #3: Treat as directed in rate chart. Thorough coverage is important. Water in granules immediately after application.

Restrictions:

- Do not apply more than six (6) times per year per location at low use rate.
- Do not apply more than three (3) times per year per location at high use rate.
- Do not allow product to run off from target site or apply within 24 hours of rain.

DIRECTIONS FOR APPLYING THIS PRODUCT

| Treatment Areas | Pests | How to Use | Use Rate |
|--|---|--|---|
| Band Treatment Around House Foundation | Ants*, Fleas, Roaches (German and American), Palmetto bugs, Water bugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs | These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses. Apply a 3 to 5 foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly immediately after application. | Apply at the rate of 0.8 lb per 1000 sq ft (an area 5' x 200') |
| Overall Lawn Application | Ants*, Armyworms (Fall & Yellow-striped), European Crane Flies, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Grasshoppers, Japanese beetle (adults), Pillbugs, Spittle bugs, Mites, Green June Beetles, Leafhoppers, Millipedes, Springtails and Ticks (including Ticks which may transmit Lyme Disease). | Apply with a fertilizer spreader. See spreader setting chart on the bag. Apply when grass is dry. Water lawn lightly immediately after application. For up to 3 month control of fire ants, apply higher use rate. | Apply at the rate of 0.8 lb per 1000 sq ft (an area 20' by 50') For control of Fire Ants apply at the rate of 2 lb per 1000 sq ft (an area 20' x 50') |
| | Billbugs, Grubs (Japanese beetle grubs, Green June Beetle Grubs), Ataenius spretulus (dung beetles), Hyperodes Weevils, Mole Crickets, Chinch bugs | Immediately water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with ¼ to ½ inch of water immediately after application. Treat again in late summer or early fall if a second generation occurs. For control of mole crickets, apply prior to or during the peak egg hatch period. | Apply 1.2 lb to 1.8 lbs per 1000 sq ft of lawn. |
| | Ants* | Water lightly immediately after application. | Apply 0.5 teaspoons over and around each ant hill. |

*Ants (Except Carpenter and Pharaoh): Argentine, Southern, Field, Allegheny Mound, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance

SPREADER SETTINGS

| Drop Spreader: | Calibration Settings at Specified Rate (lb/1000 sq ft) | | |
|--------------------------|--|-----|-----|
| | x-x | x-x | x-x |
| Scotts AccuGreen | x | x | x |
| Republic EZ #530 | x | x | x |
| Sears Drop | x | x | x |
| Rotary Spreaders: | | | |
| Scotts Deluxe | x | x | x |
| Republic EZ #540 | x | x | x |
| Earthway | x | x | x |
| Sears Rotary | x | x | x |

Note: The spreader settings listed above are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.

Treat the following pests during the months indicated. Make no more than 6 applications per year at low use rate or 3 times for highest use rate.

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|--|-----|-----|-----|-------------------|-----|-----|-----|-------|-----|-----|-----|
| | | | | Billbugs | | | | Grubs | | | |
| | | | | Dung beetle grubs | | | | | | | |
| | | | | Hyperodes weevils | | | | | | | |
| Treat Ants (except Carpenter and Pharaoh Ants) and other insects when they appear or you notice lawn damage. | | | | | | | | | | | |

STORAGE AND DISPOSAL

Storage: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.
Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.
Have the product container or label with you when calling a Poison Control Center, doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

Environmental Hazards

This product is highly toxic to fish and aquatic invertebrates. Do not apply directly to water.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Distributed by
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424
©Bonide Products, Inc.
All Rights Reserved