



ACTIVE INGREDIENTS

dicamba + 2,4-D

% ACTIVE INGREDIENTS

12.4% + 35.7%

Weedmaster contains 10.3% dicamba or 1 pound per gallon and 29.6% 2,4-D or 2.87 pounds per gallon

FORMULATION

liquid

CHEMICAL FAMILY

benzoic acids
phenoxyaliphatic acid

EPA REGISTRATION

71368-34

MODE OF ACTION

auxin growth regulator

RESTRICTED USE

no

SIGNAL WORD

danger

PACKAGING INFORMATION

2 x 2.5 gal
30 gal drum
250 gal tote
bulk

Weedmaster® contains 2,4-D and dicamba.

Weedmaster®

HERBICIDE

Depend on Weedmaster®

For years, growers and ranchers have counted on Weedmaster® for consistent weed control in sugarcane, pastures, small grains and fallow. Weedmaster is a selective, post-emerge herbicide that provides tough, long-lasting control of a broad spectrum of annual, biennial and perennial broadleaf weeds. The growth regulating effects of 2,4-D and dicamba interfere with the normal development of weeds, ultimately causing plant death.



Weedmaster[®]

HERBICIDE

LABELED USES

Conservation reserve program (CRP) land
Fallow systems
General farmstead
Sorghum
Grass (hay or silage)
Pastures
Rangeland
Sugarcane
Wheat

WEEDS CONTROLLED

ANNUALS:
Beebalm, spotted
Broomweed
Buckwheat, wild buffalobur
Burdock
Buttercup
Chickweed, common
Cockle, cow
Cocklebur, common
Coreopsis, plains
Croton, woolly
Devils-claw
Dogfennel
Evening primrose
Flax
Fleabane, annual
Flixweed
Henbit
Knotweed, spp.
Kochia
Mallow, common
Morningglory; ivyleaf, tall
Mustards; annual, tansy
Pennycress, field
Pepperweed, Virginia
Pigweed; prostrate, redroot,
smooth, tumble
Poorjoe
Purslane, common
Ragweed; common, western,
lanceleaf
Sedge
Shepherdspurse
Smartweed, Pennsylvania
Sneezeweed, bitter sowthistle
Sunflower
Thistle, Russian
Velvetleaf

BIENNIALS & PERENNIALS:
Bindweed, field
Bittercress
Buckeye species
Bullnettle*
Chicory
Clover, bur
Dandelion, common
Dewberry, southern
Dock, curly
Elderberry*
Goldenrod, Missouri
Goldenweed, common
Groundsel, Texas
Honeysuckle, hairy
Horsenettle, Carolina
Ivy, poison
Knapweed; black*,
Russian*, spotted
Lettuce, prickly
Marsheld
Mesquite
Milkweed, antelopehorn
Nightshade; silverleaf, black
Persimmon, eastern
Prickly lettuce
Rabbitbrush*
Ragwort, tansy
Redvine*
Sagebrush, fringed*
Smartweed
Sorrel, red
Sowthistle*
Spurge, leafy*
Tallow tree, Chinese
Thistle; bull, Canada*,
musk, plumeless
Vetch, hairy
Yankeeweed
Yellow

*Partial control

Key Performance Benefits

Dependable broad-spectrum control of annual and perennial broadleaves

Can improve yield or forage quality by preventing competition from weeds

Flexible application options

Excellent tank-mix partner

Application Recommendations

RATES

0.5 pints per acre to 4 - 6 pints per acre for control of annual, biennial and perennial weeds. See label for per acre rate according to weed growth stage.

Apply to actively growing weeds as aerial, broadcast, band, or spot spray applications.

Apply with water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence uses for all crops listed on this label. Post-emergence uses with sprayable fluid fertilizer may be made only on pasture, hayland, or wheat crops.

To improve burndown of emerged weeds, surfactants and/or low use rate of liquid fertilizers (28-0-0, 32-0-0), or crop oil concentrate may be used with this product or this product tank-mixes applied after the weeds have emerged.

Aerial applications: Use 3 to 10 gallons of water per acre.

Ground application (Broadcast): Use 5 to 40 gallons per broadcast acre.

FOR SPECIFIC APPLICATION RATES, DIRECTIONS, MIXING INSTRUCTIONS AND PRECAUTIONS, READ THE PRODUCT LABEL.

PLEASE VISIT WWW.NUFARM.COM/US TO DOWNLOAD A FULL LABEL.

©2009 Nufarm. Important: Always read and follow label instructions.
Weedmaster[®] is a trademark licensed to Nufarm.