

## 

Derived from calcium chloride

HELPFUL INFORMATION: Tomato blossom-end rot is a disorder caused by calcium deficiency. The disorder can be aggravated by the application of too much nitrogen fertilizer to tomato plants. Blossom-end rot appears as a round, brown sunken area on the end of the tomato fruit opposite the stem. As a fruit ripers the area turns dry and leathery. The disorder often appears after a period of rapid growth followed by dry

conditions, or in periods of heavy rain that caused calcium to leach from the soil.

DIRECTIONS FOR USE: Apply Only As Directed. Apply Bonide Rot-Stop at the rate of 2 fl. oz. (4 Tbs.) per gallon of water. Apply to the point of run-off onto the foliage and fruit during periods of rapid growth or following excessive rain fall. Apply in early morning or evening when temperatures are lower to avoid foliage burn. Make applications 5 to 7 days apart when conditions are favorable for appearance of the disorder. Regular spray programs are not recommended as prolonged spraying and rates in excess of those specified can cause foliage burn.

Foliar fertilization is intended as a supplement to a regular fertilization program and may not, by itself, provide all the nutrients normally required by crops or other intended plants. Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label, and referring to the batch number found on this container.

Information regarding the contents and levels of metals in this product is available on the internet at http://

www.aapfco.org/metals.html

STORAGE: Store & transport in an upright position. Store in the original container in an area out of reach

Buyer's Guarantee Limited to Label Claims and Refund of Purchase Price Only.

Guaranteed by: Bonide Products, Inc. F1290 6301 Sutliff Road Oriskany, NY 13424

of children & pets.

